

Comment on "Critical properties of highly frustrated pyrochlore antiferromagnets"

A. Mailhot and M.L. Plumer

Centre de Recherche en Physique du Solide et Département de Physique

Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1

(January 1993)

Abstract

We argue that the analysis of Reimers *et al.* [Phys. Rev. B **45**, 7295 (1992)] of their Monte Carlo data on the Heisenberg pyrochlore antiferromagnet, which suggests a new universality class, is not conclusive. By re-analysis of their data, we demonstrate asymptotic volume dependence in some thermodynamic quantities, which suggests the possibility that the transition may be first order.

75.40.Mg, 75.40.Cx, 75.10.-b

Reimers, Greedan and Björgvinsson¹ have recently presented results of Monte Carlo simulations of the pyrochlore Heisenberg antiferromagnet. An important issue, as they have emphasised, is to determine the order of the phase transition. This highly frustrated system is interesting from this point of view since Azaria *et al.*² have conjectured that frustration of this type can easily induce a first-order transition. Unfortunately, the determination of the order of a transition by numerical simulations on finite systems is not always straightforward, as emphasized in Ref.[1]. Peczak and Landau,³ note that if the correlation length for a system with a very weak first-order transition is much larger than the system size, then pseudocritical behavior may be observed. Scaling of the minima near T_N in Binder's fourth-order energy cumulant⁴, $V = 1 - \frac{1}{3} \frac{\langle E^4 \rangle}{\langle E^2 \rangle^2}$, with system size L (for, e.g. an $L \times L \times L$ lattice) is believed to be useful for the characterization of the order of a transition; however, a decisive conclusion based on this quantity alone can be precarious. This is due to the fact that one can never, by simulations of finite systems, determine if V_{min}^* is *exactly* $\frac{2}{3}$ or less than this value, as expected for continuous and first-order transitions, respectively. A value slightly less than $\frac{2}{3}$ may indicate either a continuous or very weakly first-order transition. Nevertheless, it is of interest to note that the 5-state Potts model in 2D, which is considered to exhibit one of the weakest first-order transitions known, has a correlation length estimated to be on the order of 2000 lattice spacings and meaningful results have been obtained from simulations of lattices that were a factor of 10 smaller⁵. The extrapolated value, $V_{min}^* \simeq 0.6655$, is consistent with expectations.

In addition to V_{min}^* , Reimers *et al.*¹ considered the scaling behavior of the specific heat and susceptibility maxima near T_N , $C_{max}(L)$ and $\chi_{max}(L)$. The pyrochlore lattice contains 16 spins per unit cell and simulations were performed on lattice sizes $L=3-10$ (with $L=9$ omitted). In terms of more familiar Bravais lattices with one spin per unit cell, the number of sites involved correspond approximately to $\tilde{L}=7.6 - 25.2$. It is certainly not clear if these lattices are large enough to reveal the true critical behavior of a highly frustrated system. Using data for *all* lattice sizes (except in the case of $V_{min}(L)$ where $L=3$ results were excluded), they found exponent ratios estimated to be $\alpha/\nu=1.8(2)$ and $1.7(2)$, from

the scaling of V_{min} and C_{max} , respectively, as well as $\gamma/\nu=3.5(3)$ from results for χ_{max} . In the case of a first order transition, these thermodynamic quantities should scale as L^3 . They also made the extrapolation $V_{min}^*=0.66658(10)$ from their data. With these values, as well as other scaling results for β and ν , they conclude that the transition is continuous and not within any known universality class.

Figs. 1, 2 and 3 present an alternative scaling analysis, namely as a function of L^3 , of their data for $\frac{2}{3}-V_{min}$, C_{max} and χ_{max} , respectively. The results suggest that the asymptotic behavior of these quantities are in fact consistent with the transition being first order. It appears possible that it is only for L larger than about 6 or 7 do the simulation results become meaningful, as indicated by the best-fit straight lines to data points for these lattice sizes. The obvious exceptions are the specific heat and energy cumulant result for $L=8$, for which we have no explanation. Our best-fit estimate for V_{min}^* from Fig. 1 is $0.66643(10)$, notably further than from the value $\frac{2}{3}$ than the result of Reimers *et al.*¹.

From this analysis, it is clear that the hypothesis of a first order transition associated with the pyrochlore antiferromagnet can be taken seriously. The results of further simulations on larger lattices would be necessary in order to make a more definitive statement regarding the critical behavior of this system. It is unlikely that experimental work can resolve this issue. Even if the transition is in fact weakly first order, there may well be a significant region of temperature near T_N where pseudocritical exponents can be measured, with results consistent with those estimated by Reimers *et al.*¹. This was indeed found to be the case for the exponent β .

This work was supported by NSERC of Canada and FCAR du Québec. We thank J. Reimers for useful discussions.

REFERENCES

- ¹ J.N. Reimers, J.E. Greedan, and M. Björgvinsson, Phys. Rev. B **45**, 7295 (1992).
- ² P. Azaria, B. Delamotte, and T. Jolicoeur, Phys. Rev. Lett. **64**, 3175 (1990); J. Appl. Phys. **69**, 6170 (1991).
- ³ P. Peczak and D.P. Landau, Phys. Rev. B. **39**, 11932 (1989).
- ⁴ M.S.S. Challa, D.P. Landau, and K. Binder, Phys. Rev. B **34**, 1841 (1986).
- ⁵ M. Fukugita, H. Mino, M. Okawa, and A. Ukawa, J. Phys. A **23**, L561 (1990).

FIGURES

FIG. 1. Scaling of the energy-cumulant minima with volume. The straight line is a fit to the data $L=6-10$

FIG. 2. Scaling of the specific-heat maxima with volume. The straight line is a fit to the data at $L=6, 7$, and 10 .

FIG. 3. Scaling of the susceptibility maxima with volume. The straight line is a fit to the data $L=7-10$